

Optimality Conditions for Nonlinear Optimization

GIAN Short Course on Optimization:
Applications, Algorithms, and Computation

Sven Leyffer

Argonne National Laboratory

September 12-24, 2016

Outline

- 1 Preliminaries: Definitions and Notation
- 2 First-Order Conditions
 - Equality Constrained Nonlinear Programs
 - Inequality Constrained Nonlinear Programs
 - The Karush-Kuhn-Tucker Conditions
- 3 Second-Order Conditions
 - Second-Order Conditions for Equality Constraints
 - Second-Order Conditions for Inequality Constraints



Preliminaries: Definitions and Notation

Seek optimality conditions for (local) minimizer ...

Definition (Nonlinear Optimization Problem)

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & c_i(x) = 0, \quad i \in \mathcal{E} \\ & l_j \leq c_i(x) \leq u_j \quad i \in \mathcal{I} \\ & l_i \leq x_i \leq u_i \quad i = 1, \dots, n \end{array}$$

where

- $f(x)$ and $c_i(x)$ twice continuously differentiable.
- \mathcal{E} indexes equality, \mathcal{I} indexes inequality constraints
- Bounds l_j, u_j, l_i, u_i can be finite or infinite

Also referred to as *nonlinear program (NLP)*.

Often, have additional structure, that can be exploited by solver



Preliminaries: Definitions and Notation

Simplify notation ... other NLPs can be expressed like this.

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & c_i(x) = 0 \quad i \in \mathcal{E} \\ & c_i(x) \geq 0 \quad i \in \mathcal{I}. \end{array}$$

Notation

$c_{\mathcal{E}}(x) = 0$, $c_{\mathcal{I}}(x) \geq 0$ denotes equality, inequality constraints.

$$\text{For } \mathcal{E} = \{1, \dots, m\} \quad c_{\mathcal{E}}(x) = \begin{pmatrix} c_1(x) \\ \vdots \\ c_m(x) \end{pmatrix}$$



Preliminaries: Definitions and Notation

Nonlinear optimization problem

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & c_i(x) = 0 \quad i \in \mathcal{E} \\ & c_i(x) \geq 0 \quad i \in \mathcal{I}. \end{array}$$

Definition (Feasible Set)

Feasible set of NLP is set of x that satisfy all constraints

$$\mathcal{F} := \left\{ x \mid c_{\mathcal{E}}(x) = 0, \text{ and } c_{\mathcal{I}}(x) \geq 0 \right\}$$

Definition (Local and Global Minimizers)

- $x^* \in \mathcal{F}$ is **global minimizer**, iff $f(x^*) \leq f(x)$ for all $x \in \mathcal{F}$.
- $x^* \in \mathcal{F}$ is **local minimizer**, iff there exists neighborhood $\mathcal{N}(x^*)$ of x^* such that $f(x^*) \leq f(x)$ for all $x \in \mathcal{F} \cup \mathcal{N}(x^*)$.

Local versus Global Minimizers

Notation.

Gradient of $f(x)$ is $g(x) = \nabla f(x)$, Jacobian of $c(x)$ is $A(x) = \nabla c^T(x)$.

Remark (Limitations of Optimality Conditions)

- *Optimality conditions only provide local optimality.*
- *Limited to smooth finite-dimensional problems.*
... extend to nonsmooth problems using subdifferential $\partial f(x)$

Remark (Importance of Optimality Conditions)

- *Guarantee that candidate solution is local optimum*
- *Indicate when point is not optimal (necessary conditions)*
- *Guide development of optimization methods*



Outline

- 1 Preliminaries: Definitions and Notation
- 2 First-Order Conditions
 - Equality Constrained Nonlinear Programs
 - Inequality Constrained Nonlinear Programs
 - The Karush-Kuhn-Tucker Conditions
- 3 Second-Order Conditions
 - Second-Order Conditions for Equality Constraints
 - Second-Order Conditions for Inequality Constraints



First-Order Conditions

Theorem (Unconstrained First-Order Conditions)

x^* unconstrained local minimizer $\Rightarrow g^* = 0$.

State this condition equivalently as

$$g^* = 0 \Leftrightarrow s^T g^* = 0, \forall s \Leftrightarrow \{s \mid s^T g^* < 0\} = \emptyset,$$

i.e. there are no strict descend directions at x^*

Generalize these conditions

- Must classify feasible directions
- Derive easy-to-check conditions for


$$\{s \mid s^T g^* < 0, \forall s \text{ feasible directions}\} = \emptyset,$$

i.e. there exist no feasible descend directions.



Concept of Feasible Directions

Feasible directions play central role in optimality ...



Distinguish two cases:

- 1 Equality constraints only.
- 2 Inequality constraints.

... equality constraints easier

Equality Constrained Nonlinear Programs

Consider equality constraints only:

$$\begin{aligned} & \text{minimize } f(x) \\ & \text{subject to } c_{\mathcal{E}}(x) = 0. \end{aligned}$$

Take infinitesimal step δ from x^* , look at Taylor series expansion:

$$c_i(x^* + \delta) = c_i(x^*) + \delta^T a_i^* + o(\|\delta\|) = \delta^T a_i^* + o(\|\delta\|),$$

because $c_i(x^*) = 0$, where $a_i^* = \nabla c_i(x^*)$

Recall: $a = o(h)$ means $\frac{a}{h} \rightarrow 0$ as $h \rightarrow 0$


Sufficient Condition for Feasible $x^* + \delta$

$$\delta^T a_i^* + o(\|\delta\|) = 0 \quad \Rightarrow \quad s^T a_i^* = 0 \quad \text{feasible directions}$$



Graphical Interpretation of Feasible Directions

Feasible directions, s such that $s^T a_i^* = 0$ are tangent directions




Feasible directions at two different points.

Graphical Interpretation of Feasible Directions

Feasible directions, s such that $s^T a_i^* = 0$ are tangent directions

How to derive feasible directions:

$$\mathcal{F} = \{x \mid x_1^2 - x_2 \leq 0, x_1^2 + x_2^2 \leq 1\}$$



$$\nabla c_1(x) = \begin{pmatrix} 2x_1 \\ -1 \end{pmatrix}$$

$$\nabla c_2(x) = \begin{pmatrix} 2x_1 \\ 2x_2 \end{pmatrix}$$

At $x = (0; 1)$ get $s = (\pm 1; 0)$:

$$(\pm 1; 0)^T (0; 2) = 0$$

At $x = (0.7861; 0.6180)$ get
two directions

$$\begin{pmatrix} -0.5367 \\ -0.8438 \end{pmatrix} \quad \begin{pmatrix} -0.6180 \\ 0.7861 \end{pmatrix}$$

Regularity Assumptions

To derive stationarity conditions, need regularity assumption:
“linearized feasible set”, looks like nonlinear feasible set

Assumption (Linear Independence of Constraint Normals)

$a_i^* = \nabla c_i(x^*)$, for $i = 1, \dots, m_e$, are linearly independent.

An alternative assumption is that all constraints are linear

- Any linearization of a linear constraint is perfect approx.
- Hence, do not need regularity assumptions for LPs and QPs.



Necessary Condition for Equality Constraints

$$\text{minimize } f(x) \quad \text{subject to } c_{\mathcal{E}}(x) = 0$$

Necessary condition: under linear independence assumption:

$$x^* \text{ is a local minimizer} \Rightarrow \left\{ s \mid s^T g^* < 0, s^T a_i^* = 0, \forall i \in \mathcal{E} \right\} = \emptyset$$

... very difficult to check

Lemma (Necessary Condition for Equality Constraints)

Assume linear independence holds, and x^ is local minimizer, then the following conditions are equivalent:*

- 1 $\left\{ s \mid s^T g^* < 0, s^T a_i^* = 0, \forall i \in \mathcal{E} \right\} = \emptyset$
- 2 *There exist Lagrange multipliers, y_i^* , for $i \in \mathcal{E}$ such that*

$$g^* = \sum_{i \in \mathcal{E}} y_i^* a_i^* = A^* y.$$

Graphic Interpretation of FO Conditions

Lemma (Necessary Condition for Equality Constraints)

Assume linear independence holds, and x^* is local minimizer, then the following conditions are equivalent:

- 1 $\{s \mid s^T g^* < 0, s^T a_i^* = 0, \forall i \in \mathcal{E}\} = \emptyset$
- 2 There exist Lagrange multipliers, y_i^* , for $i \in \mathcal{E}$ such that

$$g^* = \sum_{i \in \mathcal{E}} y_i^* a_i^* = A^* y.$$


- Can write g^* as linear combination of constraint gradients, a_i^* .
- Linear-independence Assumption implies $\text{rank}(A^*) = m_e$
i.e. A^* has full rank \Rightarrow generalized inverse, A^{*+} , exists

$$y^* = A^{*+} g^*, \quad \text{where } A^{*+} = \left(A^{*T} A^* \right)^{-1} A^{*T},$$

unique multipliers, y^* , also solve $\min \|A^* y - g^*\|_2^2$



Graphic Interpretation of FO Conditions



Method of Lagrange Multipliers

Restate conditions in Lemma as system of equations in (x, y) :

$$\begin{aligned}g(x) &= A(x) y && \text{first-order condition} \\c(x) &= 0 && \text{feasibility.}\end{aligned}$$

Define **Lagrangian function**, $\mathcal{L}(x, y) := f(x) - y^T c(x)$

Method of Lagrange Multipliers

First-order optimality conditions equivalent to

$$\nabla_x \mathcal{L}(x, y) = 0, \quad \text{and} \quad \nabla_y \mathcal{L}(x, y) = 0.$$

Can apply Newton's method to nonlinear system in (x, y)

Finding stationary points \Leftrightarrow finding stationary point of Lagrangian



Effect of Perturbations: Sensitivity Analysis

Express effect of perturbation to constraint, $c_i(x) = \epsilon$ on optimum

Let $x(\epsilon)$ and $y(\epsilon)$ denote optimal values after perturbation

$$f(x(\epsilon)) = \mathcal{L}(x(\epsilon), y(\epsilon)) = f(x(\epsilon)) + y(\epsilon)^T (c(x) - \epsilon)$$

Chain rule implies

$$\frac{df}{d\epsilon_i} = \frac{d\mathcal{L}}{d\epsilon_i} = \frac{\partial x^T}{\partial \epsilon_i} \nabla_x \mathcal{L} + \frac{\partial y^T}{\partial \epsilon_i} \nabla_y \mathcal{L} + \frac{\mathcal{L}}{\partial \epsilon_i}$$

Observe, that $\nabla_x \mathcal{L}(x, y) = 0$ and $\nabla_y \mathcal{L}(x, y) = 0$, hence

$$\frac{\mathcal{L}}{\partial \epsilon_i} = y_i \Rightarrow \frac{df}{d\epsilon_i} = y_i.$$

Sensitivity Interpretation of Multipliers

Multiplier, y_i , gives rate of change in objective to perturbation right-hand-side of constraint i .

Inequality Constrained Nonlinear Programs

Now consider both equality and inequality constraints

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & c_i(x) = 0 \quad i \in \mathcal{E} \\ & c_i(x) \geq 0 \quad i \in \mathcal{I}. \end{array}$$

only need to consider **active constraints**

$$\mathcal{A}^* := \mathcal{A}(x^*) := \{i \in \mathcal{E} \cup \mathcal{I} \mid c_i(x^*) = 0\} \quad \text{active set.}$$

... includes all equality constraints

Again, looking for feasible directions ... now for inequalities.



Inequality Constrained Nonlinear Programs

Now consider both equality and inequality constraints

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & c_i(x) = 0 \quad i \in \mathcal{E} \\ & c_i(x) \geq 0 \quad i \in \mathcal{I}. \end{array}$$

Let δ be small incremental step for active inequality, $i \in \mathcal{I} \cap \mathcal{A}^*$:

$$c_i(x^* + \delta) = c_i(x^*) + \delta^T a_i^* + o(\|\delta\|) = \delta^T a_i^* + o(\|\delta\|).$$

Now require step to remain feasible only wrt one side:

$$c_i(x^* + \delta) \geq 0 \Leftrightarrow \delta^T a_i^* + o(\|\delta\|) \geq 0$$

Hence, δ lies in direction s :

$$\text{feasible directions} \quad s^T a_i^* \geq 0, \quad \forall i \in \mathcal{I} \cap \mathcal{A}^*, \quad s^T a_i^* = 0, \quad \forall i \in \mathcal{E}.$$

... again need a regularity assumption ...



Regularity Assumption for Inequality Constraints

Need regularity assumption to ensure that linearized analysis captures nonlinear geometry

Assumption (Linear Independence Constraint Qualification)

The linear-independence constraint qualification (LICQ) holds at x^ for the NLP, iff $a_i^* = \nabla c_i(x^*)$, for $i \in \mathcal{A}^*$, are linearly independent.*

The next assumption is slightly weaker, and implies the LICQ.

Assumption (Mangasarian-Fromowitz Constraint Qualification)

The Mangasarian-Fromowitz constraint qualification (MFCQ) holds at x^ for the NLP, iff $a_i^* = \nabla c_i(x^*)$, for $i \in \mathcal{E}$, are linearly independent, and there exists $s \neq 0$ such that*

$$s^T a_i^* > 0, \forall i \in \mathcal{I} \cap \mathcal{A}^*.$$



Why We Need Regularity Assumptions

Consider the NLP


$$\begin{array}{ll} \underset{x}{\text{minimize}} & x_1 \\ \text{subject to} & x_2 \leq x_1^3 \\ & x_2 \geq 0 \end{array}$$

Has optimum at cusp

$$x^* = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

... but constraints violate MFCQ
⇒ bogus “feasible” direction

$$s = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$$



MFCQ fails at cusp, $x = 0$

Regularity Assumption for Inequality Constraints

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & c_i(x) = 0 \quad i \in \mathcal{E} \\ & c_i(x) \geq 0 \quad i \in \mathcal{I}. \end{array}$$

Assumption (Mangasarian-Fromowitz Constraint Qualification)

The Mangasarian-Fromowitz constraint qualification (MFCQ) holds at x^ for the NLP, iff $a_i^* = \nabla c_i(x^*)$, for $i \in \mathcal{E}$, are linearly independent, and there exists $s \neq 0$ such that*

$$s^T a_i^* > 0, \forall i \in \mathcal{I} \cap \mathcal{A}^*.$$

MFCQ is stronger than needed:

$$\left\{ s \mid s^T g^* < 0, s^T a_i^* = 0, \forall i \in \mathcal{E}, s^T a_i^* \geq 0, \forall i \in \mathcal{I} \cap \mathcal{A}^* \right\} = \emptyset$$

... but this condition really difficult to check.



Necessary Condition for Nonlinear Optimization

Lemma (First-Order Conditions for Optimality)

Assume that LICQ or MFCQ hold, and that x^* is local minimizer, then the following two conditions are equivalent:

- 1 There exist no feasible descend direction:

$$\left\{ s \mid s^T g^* < 0, s^T a_i^* = 0, \forall i \in \mathcal{E}, s^T a_i^* \geq 0, \forall i \in \mathcal{I} \cap \mathcal{A}^* \right\} = \emptyset$$

- 2 There exist so-called *Lagrange multipliers*, y_i^* , for $i \in \mathcal{A}^*$:

$$g^* = \sum_{i \in \mathcal{A}^*} y_i^* a_i^* = A^* y \quad \text{where } y_i^* \geq 0, \forall i \in \mathcal{I} \cap \mathcal{A}^*.$$



Necessary Condition for Nonlinear Optimization

$$g^* = \sum_{i \in \mathcal{A}^*} y_i^* a_i^* = A^* y \quad \text{where } y_i^* \geq 0, \forall i \in \mathcal{I} \cap \mathcal{A}^*.$$

Remark (Towards an Algorithms for NLP)

Assume at non-stationary point with

- *Multiplier $\lambda_q < 0$ for some $q \in \mathcal{I}$
e.g. least-squares multiplier*
- *Have direction s with $s^T a_q = 1$*

Then reduce objective by step in this feasible direction s .

Basis for active-set methods for linear and quadratic programming!



The Karush-Kuhn-Tucker Conditions

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & c_i(x) = 0 \quad i \in \mathcal{E} \\ & c_i(x) \geq 0 \quad i \in \mathcal{I}. \end{array}$$

Theorem (Karush-Kuhn-Tucker (KKT) Conditions)

x^* local minimizer of NLP and assume LICQ or MFCQ hold at x^* .
Then there exist Lagrange multipliers, y^* such that

$$\nabla_x \mathcal{L}(x^*, y^*) = 0 \quad \text{first order condition} \quad (1)$$

$$c_{\mathcal{E}}(x^*) = 0 \quad \text{feasibility} \quad (2)$$

$$c_{\mathcal{I}}(x^*) \geq 0 \quad \text{feasibility} \quad (3)$$

$$y_{\mathcal{I}}^* \geq 0 \quad \text{dual feasibility} \quad (4)$$

$$y_i^* c_i(x^*) = 0 \quad \text{complementary slackness.} \quad (5)$$



Interpretation of KKT Conditions

Remark (Stationarity Conditions and Algorithms)

Take standard NLP & linearize about stationary point, x^* , then:
KKT conditions are the FO conditions of linearized problem:

$$\begin{aligned} & \underset{d}{\text{minimize}} && f(x^*) + d^T \nabla f(x^*) \\ & \text{subject to} && c_i(x^*) + d^T \nabla c_i(x^*) = 0, \quad i \in \mathcal{E} \\ & && c_i(x^*) + d^T \nabla c_i(x^*) \geq 0, \quad i \in \mathcal{I}, \end{aligned}$$

- Motivates **algorithms** such as SLP, SQP, SLQP, SQQP, ...
- Extends FO conditions to **structured** NLP, e.g. MPECs, ...
... and hence defines new **structured algorithmic approaches**



Outline

- 1 Preliminaries: Definitions and Notation
- 2 First-Order Conditions
 - Equality Constrained Nonlinear Programs
 - Inequality Constrained Nonlinear Programs
 - The Karush-Kuhn-Tucker Conditions
- 3 Second-Order Conditions
 - Second-Order Conditions for Equality Constraints
 - Second-Order Conditions for Inequality Constraints



Second-Order Conditions

KKT conditions are first-order necessary conditions.

Goal

Extend second-order from the unconstrained case

Remark

Important to include second-order effects from constraints

- *Can replace objective:*

$$\underset{x}{\text{minimize}} f(x) \quad \Leftrightarrow \quad \underset{x, \eta}{\text{minimize}} \eta \quad \text{subject to } \eta \geq f(x)$$

- *Need to consider $\nabla^2 c_i(x)$, not just $\nabla^2 f(x)$.*

Again convenient to distinguish equality and inequality constraints.



Second-Order Conditions for Equality Constraints

Let x^* is KKT point, and a_i^* for $i \in \mathcal{E}$ linearly independent

Let δ be an incremental step along feasible direction, s .

$$\begin{aligned} f(x^* + \delta) &= \mathcal{L}(x^* + \delta, y^*) \\ &= \mathcal{L}(x^*, y^*) + \delta^T \nabla_x \mathcal{L}(x^*, y^*) + \frac{1}{2} \delta^T W^* \delta + o(\|\delta\|^2) \\ &= f(x^*) + \frac{1}{2} \delta^T W^* \delta + o(\|\delta\|^2), \end{aligned}$$

where Hessian of Lagrangian is:

$$W^* = \nabla^2 \mathcal{L}(x^*, y^*) = \nabla^2 f(x^*) + \sum_{i \in \mathcal{E}} y_i^* \nabla^2 c_i(x^*)$$

Optimality of x^* implies

$$s^T W^* s \geq 0, \quad \forall s : s^T \nabla a_i^* = 0.$$

i.e. Lagrangian has nonnegative curvature for all feasible directions



Second-Order Conditions for Equality Constraints

Proposition (Second-Order Necessary Condition)

x^* local minimizer, and if constraint qualification holds, then

$$s^T \nabla^2 \mathcal{L}(x^*, y^*) s \geq 0, \quad \forall s : s^T \nabla a_i^* = 0.$$

Can also state sufficient condition for local minimizer.

Proposition (Second-Order Sufficient Condition)

If $\nabla_x \mathcal{L}(x^*, y^*) = 0$, if $c(x^*) = 0$, and if

$$s^T \nabla^2 \mathcal{L}(x^*, y^*) s > 0, \quad \forall s \neq 0 : s^T \nabla a_i^* = 0,$$

then x^* is a local minimizer.

Note: \exists gap between necessary and sufficient conditions.



Second-Order Conditions for Inequality Constraints

To derive second-order conditions consider active constraints, \mathcal{A}^* .
 \Rightarrow NLP equivalent to equality NLP, if $y_i^* > 0, \forall i \in \mathcal{I} \cap \mathcal{A}^*$,

Simplifying Assumption

Assume strict complementarity: $y_i^* > 0, \forall i \in \mathcal{I} \cap \mathcal{A}^*$,

Proposition (Second-Order Sufficient Condition)

If $\nabla_x \mathcal{L}(x^, y^*) = 0$, if $c(x^*) = 0$, if strict complementarity holds, i.e. $y_i^* > 0, \forall i \in \mathcal{I} \cap \mathcal{A}^*$, and if*

$$s^T \nabla^2 \mathcal{L}(x^*, y^*) s > 0, \forall s \neq 0 : s^T \nabla a_i^* = 0, \forall i \in \mathcal{A}^*,$$

then it follows that x^ is a local minimizer.*



Second-Order Conditions for Inequality Constraints

More rigorous results without strict complementarity possible ...
... needs Hessian $\nabla^2 \mathcal{L}$ positive definite over cone **impractical**

Check sufficient conditions by finding inertia of KKT matrix,

$$\begin{bmatrix} W^* & A^* \\ A^{*T} & 0 \end{bmatrix}.$$

Theorem


If inertia of KKT matrix is $[n - m, 0, m]$, then second order conditions are satisfied, where $m = |\mathcal{A}^|$.*

KKT matrix with inertia is $[n - m, 0, m]$ is **second-order sufficient**

Matrix inertia: triple of positive, zero, and negative eigenvalues.



Summary and Take-Aways



Derived Optimality Conditions for NLPs

- Intuitive geometric interpretation
- Motivate algorithmic approaches (soon)

Optimality Conditions Require Regularity

- Not easy to check a priori (LICQ is OK)
- **What happens if regularity does not hold?**
- Algorithms often detect lack of regularity ... fail “gracefully” ...

